

IGC-PFUR-08/2003

# ON COMPOSITE S-BRANE SOLUTIONS WITH ORTHOGONAL INTERSECTION RULES

**V.D. Ivashchuk<sup>1</sup>**

*Center for Gravitation and Fundamental Metrology, VNIIMS, 3/1 M.  
Ulyanovoy Str., Moscow 119313, Russia  
Institute of Gravitation and Cosmology, Peoples' Friendship University of  
Russia, 6 Miklukho-Maklaya St., Moscow 117198, Russia*

## Abstract

It is shown that non-composite intersecting  $S$ -brane solutions received recently by N. Ohta [Phys. Lett. B 558, 213 (2003); hep-th/0301095] are special case of cosmological-type solutions with composite  $p$ -branes obtained in our earlier publications.

---

<sup>1</sup>ivashchuk@mail.ru

# 1 Introduction

This paper is comment on a recent publication of N. Ohta [1] devoted to intersection rules for  $S$ -branes. (The notion “ $S$ -brane” appeared in [2], see also [3, 5] and references therein.)

The solution considered in [1] is not a new one but: (i) for non-zero value of Robertson-Walker parameter  $\sigma$  it is a special case of our solution from [4], (ii) for zero value of  $\sigma$  it is a special case of the solutions obtained earlier in [5] (see also [7]).

We note that the intersection rules considered in [1] are not new ones but they special case of so-called ”orthogonal” intersection rules from [4]. (For earlier publications see also [8, 9, 10, 11] and refs. in [1]).

Here we deal with a model governed by the action

$$S_g = \int d^D x \sqrt{|g|} \left\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \Delta} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)^2 \right\} \quad (1.1)$$

where  $g = g_{MN}(x) dx^M \otimes dx^N$  is a metric,  $\varphi = (\varphi^\alpha) \in \mathbf{R}^l$  is a vector of scalar fields,  $(h_{\alpha\beta})$  is a constant symmetric non-degenerate  $l \times l$  matrix ( $l \in \mathbf{N}$ ),  $\theta_a = \pm 1$ ,  $F^a = dA^a = \frac{1}{n_a!} F_{M_1 \dots M_{n_a}}^a dz^{M_1} \wedge \dots \wedge dz^{M_{n_a}}$  is a  $n_a$ -form ( $n_a \geq 1$ ),  $\lambda_a$  is a 1-form on  $\mathbf{R}^l$ :  $\lambda_a(\varphi) = \lambda_{\alpha a} \varphi^\alpha$ ,  $a \in \Delta$ ,  $\alpha = 1, \dots, l$ . In (1.1) we denote  $|g| = |\det(g_{MN})|$ ,  $(F^a)_g^2 = F_{M_1 \dots M_{n_a}}^a F_{N_1 \dots N_{n_a}}^a g^{M_1 N_1} \dots g^{M_{n_a} N_{n_a}}$ ,  $a \in \Delta$ . Here  $\Delta$  is some finite set. For pseudo-Euclidean metric of signature  $(-, +, \dots, +)$  all  $\theta_a = 1$ .

The paper is organized as following. In Section 2 we overview cosmological-type solutions with composite intersecting  $p$ -branes from [4, 5] obeying to ”orthogonal” intersection rules. Section 3 is devoted to scalar-vacuum parameterization of  $S$ -brane solutions that in a certain special case appears in the paper of N. Ohta [1].

## 2 Cosmological-type solutions with composite intersecting $p$ -branes

### 2.1 Solutions with $n$ Ricci-flat spaces

Let us consider a family of solutions to field equations corresponding to the action (1.1) and depending upon one variable  $u$  [5] (see also [6, 7]).

These solutions are defined on the manifold

$$M = (u_-, u_+) \times M_1 \times M_2 \times \dots \times M_n, \quad (2.1)$$

where  $(u_-, u_+)$  is an interval belonging to  $\mathbf{R}$ , and have the following form

$$g = \left( \prod_{s \in S} [f_s(u)]^{2d(I_s)h_s/(D-2)} \right) \left\{ \exp(2c^0 u + 2\bar{c}^0) w du \otimes du + \right. \quad (2.2)$$

$$\left. \sum_{i=1}^n \left( \prod_{s \in S} [f_s(u)]^{-2h_s \delta_{iI_s}} \right) \exp(2c^i u + 2\bar{c}^i) g^i \right\},$$

$$\exp(\varphi^\alpha) = \left( \prod_{s \in S} f_s^{h_s \chi_s \lambda_{a_s}^\alpha} \right) \exp(c^\alpha u + \bar{c}^\alpha), \quad (2.3)$$

$$F^a = \sum_{s \in S} \delta_{a_s}^a \mathcal{F}^s, \quad (2.4)$$

$\alpha = 1, \dots, l; a \in \Delta$ .

In (2.2)  $w = \pm 1$ ,  $g^i = g_{m_i n_i}^i(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$  is a Ricci-flat metric on  $M_i$ ,  $i = 1, \dots, n$ ,

$$\delta_{iI} = \sum_{j \in I} \delta_{ij} \quad (2.5)$$

is the indicator of  $i$  belonging to  $I$ :  $\delta_{iI} = 1$  for  $i \in I$  and  $\delta_{iI} = 0$  otherwise.

The  $p$ -brane set  $S$  is by definition

$$S = S_e \sqcup S_m, \quad S_v = \sqcup_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \quad (2.6)$$

$v = e, m$  and  $\Omega_{a,e}, \Omega_{a,m} \subset \Omega$ , where  $\Omega = \Omega(n)$  is the set of all non-empty subsets of  $\{1, \dots, n\}$ . Here and in what follows  $\sqcup$  means the union of non-intersecting sets. Any  $p$ -brane index  $s \in S$  has the form

$$s = (a_s, v_s, I_s), \quad (2.7)$$

where  $a_s \in \Delta$  is colour index,  $v_s = e, m$  is electro-magnetic index and the set  $I_s \in \Omega_{a_s, v_s}$  describes the location of  $p$ -brane worldvolume.

The sets  $S_e$  and  $S_m$  define electric and magnetic  $p$ -branes, correspondingly. In (2.3)

$$\chi_s = +1, -1 \quad (2.8)$$

for  $s \in S_e, S_m$ , respectively. In (2.4) forms

$$\mathcal{F}^s = Q_s f_s^{-2} du \wedge \tau(I_s), \quad (2.9)$$

$s \in S_e$ , correspond to electric  $p$ -branes and forms

$$\mathcal{F}^s = Q_s \tau(\bar{I}_s), \quad (2.10)$$

$s \in S_m$ , correspond to magnetic  $p$ -branes;  $Q_s \neq 0$ ,  $s \in S$ . Here and in what follows

$$\bar{I} \equiv I_0 \setminus I, \quad I_0 = \{1, \dots, n\}. \quad (2.11)$$

All manifolds  $M_i$  are assumed to be oriented and connected and the volume  $d_i$ -forms

$$\tau_i \equiv \sqrt{|g^i(y_i)|} dy_i^1 \wedge \dots \wedge dy_i^{d_i}, \quad (2.12)$$

and parameters

$$\varepsilon(i) \equiv \text{sign}(\det(g_{m_i n_i}^i)) = \pm 1 \quad (2.13)$$

are well-defined for all  $i = 1, \dots, n$ . Here  $d_i = \dim M_i$ ,  $i = 1, \dots, n$ ;  $D = 1 + \sum_{i=1}^n d_i$ . For any set  $I = \{i_1, \dots, i_k\} \in \Omega$ ,  $i_1 < \dots < i_k$ , we denote

$$\tau(I) \equiv \tau_{i_1} \wedge \dots \wedge \tau_{i_k}, \quad (2.14)$$

$$d(I) \equiv \sum_{i \in I} d_i, \quad (2.15)$$

$$\varepsilon(I) \equiv \varepsilon(i_1) \dots \varepsilon(i_k). \quad (2.16)$$

The parameters  $h_s$  appearing in the solution satisfy the relations

$$h_s = (B_{ss})^{-1}, \quad (2.17)$$

where

$$B_{ss'} \equiv d(I_s \cap I_{s'}) + \frac{d(I_s)d(I_{s'})}{2-D} + \chi_s \chi_{s'} \lambda_{\alpha a_s} \lambda_{\beta a_{s'}} h^{\alpha\beta}, \quad (2.18)$$

$s, s' \in S$ , with  $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$ .

Here we assume that

$$(i) \quad B_{ss} \neq 0, \quad (2.19)$$

for all  $s \in S$ , and

$$(ii) \quad B_{ss'} = 0, \quad (2.20)$$

for  $s \neq s'$ , i.e. canonical (orthogonal) intersection rules are satisfied.

The moduli functions read

$$f_s(u) = R_s \sinh(\sqrt{C_s}(u - u_s)), \quad C_s > 0, \quad h_s \varepsilon_s < 0; \quad (2.21)$$

$$R_s \sin(\sqrt{|C_s|}(u - u_s)), \quad C_s < 0, \quad h_s \varepsilon_s < 0; \quad (2.22)$$

$$R_s \cosh(\sqrt{C_s}(u - u_s)), \quad C_s > 0, \quad h_s \varepsilon_s > 0; \quad (2.23)$$

$$|Q^s| |h_s|^{-1/2} (u - u_s), \quad C_s = 0, \quad h_s \varepsilon_s < 0, \quad (2.24)$$

where  $R_s = |Q_s| |h_s C_s|^{-1/2}$ ,  $C_s$ ,  $u_s$  are constants,  $s \in S$ .

Here

$$\varepsilon_s = (-\varepsilon[g])^{(1-\chi_s)/2} \varepsilon(I_s) \theta_{a_s}, \quad (2.25)$$

$s \in S$ ,  $\varepsilon[g] \equiv \text{sign}(\det(g_{MN}))$ . More explicitly (2.25) reads:  $\varepsilon_s = \varepsilon(I_s) \theta_{a_s}$  for  $v_s = e$  and  $\varepsilon_s = -\varepsilon[g] \varepsilon(I_s) \theta_{a_s}$  for  $v_s = m$ .

Vectors  $c = (c^A) = (c^i, c^\alpha)$  and  $\bar{c} = (\bar{c}^A)$  obey the following constraints

$$\sum_{i \in I_s} d_i c^i - \chi_s \lambda_{a_s \alpha} c^\alpha = 0, \quad \sum_{i \in I_s} d_i \bar{c}^i - \chi_s \lambda_{a_s \alpha} \bar{c}^\alpha = 0, \quad s \in S, \quad (2.26)$$

$$c^0 = \sum_{j=1}^n d_j c^j, \quad \bar{c}^0 = \sum_{j=1}^n d_j \bar{c}^j, \quad (2.27)$$

$$\sum_{s \in S} C_s h_s + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^n d_i (c^i)^2 - \left( \sum_{i=1}^n d_i c^i \right)^2 = 0. \quad (2.28)$$

Here we identify notations for  $g^i$  and  $\hat{g}^i$ , where  $\hat{g}^i = p_i^* g^i$  is the pullback of the metric  $g^i$  to the manifold  $M$  by the canonical projection:  $p_i : M \rightarrow M_i$ ,  $i = 1, \dots, n$ . An analogous agreement will be also kept for volume forms etc.

Due to (2.9) and (2.10), the dimension of  $p$ -brane worldvolume  $d(I_s)$  is defined by

$$d(I_s) = n_{a_s} - 1, \quad d(I_s) = D - n_{a_s} - 1, \quad (2.29)$$

for  $s \in S_e, S_m$ , respectively. For a  $p$ -brane we have  $p = p_s = d(I_s) - 1$ .

## 2.2 Solutions with one curved Einstein space and $n-1$ Ricci-flat spaces

The cosmological solution with Ricci-flat spaces may be also modified to the following case:

$$\text{Ric}[g^1] = \xi_1 g^1, \quad \xi_1 \neq 0, \quad \text{Ric}[g^i] = 0, \quad i > 1, \quad (2.30)$$

i.e. the first space  $(M_1, g^1)$  is Einstein space of non-zero scalar curvature and other spaces  $(M_i, g^i)$  are Ricci-flat and

$$1 \notin I_s, \quad \forall s \in S, \quad (2.31)$$

i.e. all “brane” sub-manifolds do not contain  $M_1$ .

In this case the exact solution may be obtained by a little modifications of the solutions from the previous subsection. The metric reads as follows [4]

$$g = \left( \prod_{s \in S} [f_s(u)]^{2d(I_s)h_s/(D-2)} \right) \left\{ [f_1(u)]^{2d_1/(1-d_1)} \exp(2c^1 u + 2\bar{c}^1) \times \right. \quad (2.32)$$

$$\left. \times [w du \otimes du + f_1^2(u) g^1] + \sum_{i=2}^n \left( \prod_{s \in S} [f_s(u)]^{-2h_s \delta_{iI_s}} \right) \exp(2c^i u + 2\bar{c}^i) g^i \right\}.$$

where

$$f_1(u) = R \sinh(\sqrt{C_1}(u - u_1)), \quad C_1 > 0, \quad \xi_1 w > 0; \quad (2.33)$$

$$R \sin(\sqrt{|C_1|}(u - u_1)), \quad C_1 < 0, \quad \xi_1 w > 0; \quad (2.34)$$

$$R \cosh(\sqrt{C_1}(u - u_1)), \quad C_1 > 0, \quad \xi_1 w < 0; \quad (2.35)$$

$$|\xi_1(d_1 - 1)|^{1/2} (u - u_1), \quad C_1 = 0, \quad \xi_1 w > 0, \quad (2.36)$$

$u_1$  and  $C_1$  are constants,  $R = |\xi_1(d_1 - 1)/C_1|^{1/2}$ , and

$$\sum_{s \in S} C_s h_s + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^n d_i (c^i)^2 - \left( \sum_{i=1}^n d_i c^i \right)^2 = C_1 \frac{d_1}{d_1 - 1}. \quad (2.37)$$

Now, vectors  $c = (c^A)$  and  $\bar{c} = (\bar{c}^A)$  satisfy also additional constraints

$$c^1 = \sum_{j=1}^n d_j c^j, \quad \bar{c}^1 = \sum_{j=1}^n d_j \bar{c}^j. \quad (2.38)$$

All other relations from the previous subsection are unchanged.

**Restrictions on  $p$ -brane configurations.** The solutions presented above are valid if two restrictions on the sets of composite  $p$ -branes are satisfied [6]. These restrictions guarantee the block-diagonal form of the energy-momentum tensor and the existence of the sigma-model representation (without additional constraints) [11].

The first restriction reads

$$(\mathbf{R1}) \quad d(I \cap J) \leq d(I) - 2, \quad (2.39)$$

for any  $I, J \in \Omega_{a,v}$ ,  $a \in \Delta$ ,  $v = e, m$  (here  $d(I) = d(J)$ ).

The second restriction is following one

$$(\mathbf{R2}) \quad d(I \cap J) \neq 0, \quad (2.40)$$

for  $I \in \Omega_{a,e}$  and  $J \in \Omega_{a,m}$ ,  $a \in \Delta$ .

### 3 Scalar-vacuum parameterization of $p$ -brane solutions

Here we show that the solutions with  $p$ -branes may be written in a parameterization that appears in the absence of  $p$ -branes (when  $Q_s = 0$ ).

#### 3.1 Scalar-vacuum solutions

Let us consider the scalar vacuum case when all charges  $Q_s$  are zero.

##### 3.1.1 Solution with Ricci-flat spaces

The scalar vacuum analogue of the solution from subsection 2.1 reads

$$g = \exp(2c^0 u + 2\bar{c}^0) w du \otimes du + \sum_{i=1}^n \exp(2c^i u + 2\bar{c}^i) g^i, \quad (3.1)$$

$$\varphi^\alpha = c^\alpha u + \bar{c}^\alpha, \quad (3.2)$$

$\alpha = 1, \dots, l$ ; where  $c^0$  and  $\bar{c}^0$  obey to (2.27) and

$$h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^n d_i (c^i)^2 - \left( \sum_{i=1}^n d_i c^i \right)^2 = 0. \quad (3.3)$$

A special cases of this Kasner-like solution were considered in [12] (without scalar fields) and in [13] (with one scalar field).

##### 3.1.2 Solutions with one Einstein space of non-zero curvature

The scalar vacuum analogue of the solution from subsection 2.2 is the following one

$$g = \exp(2c^1 u + 2\bar{c}^1) [w du \otimes du + f_1^2(u) g^1] + \sum_{i=2}^n \exp(2c^i u + 2\bar{c}^i) g^i, \quad (3.4)$$

where  $f_1(u)$  is defined in (2.33)-(2.36) and

$$h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^n d_i (c^i)^2 - \left( \sum_{i=1}^n d_i c^i \right)^2 = C_1 \frac{d_1}{d_1 - 1}. \quad (3.5)$$

Here  $c^1$  and  $\bar{c}^1$  obey to linear constraints (2.38).

Special cases of this solution were considered in [12] (without scalar fields) and in [14, 16, 15] (for one scalar field).

### 3.2 Minisuperspace-covariant notations

Here we show that the constraint (2.37) may be rewritten in a scalar-vacuum form (3.5). To do this we use the following minisuperspace-covariant relations [4, 11]:

$$(\bar{G}_{AB}) = \begin{pmatrix} G_{ij} & 0 \\ 0 & h_{\alpha\beta} \end{pmatrix}, \quad (\bar{G}^{AB}) = \begin{pmatrix} G^{ij} & 0 \\ 0 & h^{\alpha\beta} \end{pmatrix} \quad (3.6)$$

are, correspondingly, a (truncated) target space metric and inverse to it, where (see [17])

$$G_{ij} = d_i \delta_{ij} - d_i d_j, \quad G^{ij} = \frac{\delta^{ij}}{d_i} + \frac{1}{2-D}, \quad (3.7)$$

and

$$U_A^s c^A = \sum_{i \in I_s} d_i c^i - \chi_s \lambda_{a_s \alpha} c^\alpha, \quad (U_A^s) = (d_i \delta_{i I_s}, -\chi_s \lambda_{a_s \alpha}), \quad (3.8)$$

are co-vectors,  $s = (a_s, v_s, I_s) \in S$  and  $(c^A) = (c^i, c^\alpha)$ .

In what follows we use a scalar product [11]

$$(U, U') = \bar{G}^{AB} U_A U'_B, \quad (3.9)$$

for  $U = (U_A), U' = (U'_A) \in \mathbf{R}^N$ ,  $N = n + l$ .

The scalar products for vectors  $U^s$  were calculated in [11]

$$(U^s, U^{s'}) = B_{ss'}, \quad (3.10)$$

where  $s = (a_s, v_s, I_s)$ ,  $s' = (a_{s'}, v_{s'}, I_{s'})$  belong to  $S$  and  $B_{ss'}$  are defined in (2.18). Due to relations (2.20)  $U^s$ -vectors are orthogonal, i.e.

$$(U^s, U^{s'}) = 0, \quad (3.11)$$

for  $s \neq s'$ .

The linear and quadratic constraints from (2.26), (2.38) and (2.37), respectively, read in minisuperspace-covariant form as follows:

$$U_A^s c^A = 0, \quad U_A^s \bar{c}^A = 0, \quad (3.12)$$

$s \in S$ ,

$$U_A^1 c^A = 0, \quad U_A^1 \bar{c}^A = 0, \quad (3.13)$$

and

$$\sum_{s \in S} C_s h_s + \bar{G}_{AB} c^A c^B = C_1 \frac{d_1}{d_1 - 1}. \quad (3.14)$$



In (3.13)

$$U_A^1 c^A = -c^1 + d_i c^i, \quad (U_A^1) = (-\delta_i^1 + d_i, 0), \quad (3.15)$$

is a co-vector corresponding to curvature term. It obeys

$$(U^1, U^1) = \frac{1}{d_1} - 1 < 0, \quad (U^1, U^s) = 0 \quad (3.16)$$

for all  $s \in S$ . The last relation (3.16) follows from (2.31).

### 3.3 New parameters for $S$ -branes

Let us consider the special case of the solutions from sect. 2.2 with  $h_s \varepsilon_s > 0$  for all  $s \in S$  and  $w = -1$ . When all  $\theta_a > 0$  and all internal spaces are of Euclidean signature (and hence all  $\varepsilon_s > 0$ ) we are led to  $S$ -brane solutions (with  $h_s$ ) [5].

We get from (2.21)-(2.24) that in this case

$$f_s = R_s \cosh(\sqrt{C_s}(u - u_s)), \quad (3.17)$$

where  $R_s = |Q_s| |h_s C_s|^{-1/2}$  and  $C_s > 0$  for all  $s \in S$ .

Let us introduce new parameters

$$c_{new}^A \equiv c^A - \sum_{s \in S} h_s \sqrt{C_s} U^{sA}, \quad (3.18)$$

$$\bar{c}_{new}^A \equiv \bar{c}^A - \sum_{s \in S} h_s \sqrt{C_s} U^{sA}, \quad (3.19)$$

where  $U^{sA} = \bar{G}^{AB} U_B^s$ .

In terms of new parameters the relations (3.12)-(3.14) read

$$U_A^s c_{new}^A = U_A^s \bar{c}_{new}^A = -\sqrt{C_s}, \quad (3.20)$$

$$U_A^1 c_{new}^A = U_A^1 \bar{c}_{new}^A = 0, \quad (3.21)$$

$$\bar{G}_{AB} c_{new}^A c_{new}^B = C_1 \frac{d_1}{d_1 - 1}. \quad (3.22)$$

The last relation when written in components has a scalar-vacuum form (3.5), or using relations (3.21) (see (3.15)) we get

$$h_{\alpha\beta} c_{new}^\alpha c_{new}^\beta + \sum_{i=2}^n d_i (c_{new}^i)^2 + \frac{1}{d_1 - 1} \left( \sum_{i=2}^n d_i c_{new}^i \right)^2 = C_1 \frac{d_1}{d_1 - 1}. \quad (3.23)$$

In the special case of non-composite  $S$ -brane solutions with one scalar field and internal one-dimensional spaces of Euclidean signature ( $M_i$ ,  $i > 1$ ) the relation (3.23) is coinciding (up to notations) with the formula (26) from Ohta's paper [1]. In this case relations (3.17), (3.20) are in agreement with formulas (37) and (38), respectively, from [1].

Thus, in the special case under consideration when the factor space  $M_1$  is a spherical or a hyperbolic space we are led to the solution from [1].<sup>2</sup>

An analogous consideration may be carried out for solutions from sect. 2.1 when  $M_1$  is a flat space and  $S$ -branes do not "live" in  $M_1$  (see section 3 from [5]). In this case one can also verify that Ohta's solution is a special case of our solution from sect. 2.1.

## 4 Conclusions

In this paper we overviewed cosmological-type solutions with composite intersecting  $p$ -branes from [4, 5] obeying to "orthogonal" intersection rules. We obtained a scalar-vacuum parameterization of  $S$ -brane solutions that allow us to identify recent solution obtained by N. Ohta in [1] as a special case of our cosmological type (e.g.  $S$ -brane) solutions.

It should be noted that the method of derivation of solutions used by N. Ohta in [1] is not coinciding with our approach based on reduction to Coda-like systems [4, 7]. It should be also noted that our approach gives a rather systematic way to derivation of intersection rules using the integrability conditions for Coda-like systems [4, 6, 7]. In this approach intersection rules have a minisuperspace-covariant form, i.e. they are formulated in terms of scalar products of brane  $U$ -vectors and are classified by Caftan matrices of (semi-simple) Lie algebras. The intersection rules considered in this paper correspond to the Lie algebra  $A_1 + \dots + A_1$ .

## Acknowledgments

This work was supported in part by the Russian Ministry of Science and Technology, Russian Foundation for Basic Research (RFFI-01-02-17312-a) and Project DFG (436 RUS 113/678/0-1(R)).

---

<sup>2</sup>There are also two extra parameters in the solution from [1] standing for rescaling of time and  $M_1$  manifolds, respectively.

## References

- [1] N. Ohta, Intersection rules for S-branes, *Phys. Lett. B* **558**, 213 (2003); hep-th/0301095
- [2] M. Gutperle and A. Strominger, Spacelike branes, *JHEP* **0204**, 018 (2002); hep-th/0202210.
- [3] C.M. Chen, D.M. Gal'tsov and M. Gutperle, S-brane solutions in supergravity theories; *Phys. Rev. D* **66**, 024043 (2002); hep-th/0204071.
- [4] V.D. Ivashchuk and V.N. Melnikov, Multidimensional classical and quantum cosmology with intersecting  $p$ -branes, hep-th/9708157; *J. Math. Phys.*, **39**, 2866-2889 (1998).
- [5] V.D. Ivashchuk, Composite S-brane solutions related to Toda-type systems, *Class. Quantum Grav.* **20**, 261-276 (2003); hep-th/0208101.
- [6] V.D. Ivashchuk and S.-W. Kim, Solutions with intersecting p-branes related to Toda chains, *J. Math. Phys.*, **41** (1) 444-460 (2000); hep-th/9907019.
- [7] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.* **18**, R82-R157 (2001); hep-th/0110274.
- [8] V.D. Ivashchuk and V.N. Melnikov, Intersecting p-Brane Solutions in Multidimensional Gravity and M-Theory, hep-th/9612089; *Grav. and Cosmol.* **2**, No 4, 297-305 (1996); Generalized Intersecting P-brane Solutions from Sigma-model, *Phys. Lett. B* **403**, 23-30 (1997).
- [9] I.Ya. Aref'eva and O.A. Rytchkov, Incidence Matrix Description of Intersecting p-brane Solutions, hep-th/9612236.
- [10] I.Ya. Aref'eva, M.G. Ivanov and I.V. Volovich, Non-extremal intersecting p-branes in various dimensions, hep-th/9702079; *Phys. Lett. B* **406**, 44-48 (1997).
- [11] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the Generalized Composite p-branes, hep-th/9705036; *Class. Quantum Grav.* **14**, 3001-3029 (1997); Corrigenda *ibid.* **15** (12), 3941 (1998).
- [12] V.D. Ivashchuk, Multidimensional Cosmology and Toda-like Systems, *Phys. Lett. , A* **170**, 16-20 (1992).
- [13] U. Bleyer and A. Zhuk, Kasner-like, inflationary and steady-state solutions in multidimensional cosmology, *Astron. Nachrichten*, **317**, 161-173 (1996).
- [14] U. Bleyer and A. Zhuk, Multidimensional integrable cosmological models with negative external curvature, *Grav. Cosmol.*, **1**, 106-118 (1995).
- [15] U. Bleyer and A. Zhuk, Multidimensional integrable cosmological models with positive external space curvature, *Grav. Cosmol.* **1**, 37-45 (1995).

- [16] V.D. Ivashchuk and V.N.Melnikov, Multidimensional classical and quantum cosmology with perfect fluid, *Grav. Cosmol.*, **1**, 133-148 (1995); hep-th/9503223.
- [17] V.D. Ivashchuk, V.N. Melnikov and A.I. Zhuk, On Wheeler-DeWitt Equation in Multidimensional Cosmology, *Nuovo Cimento* **B 104**, 575 (1989).